



This open access document is published as a preprint in the Beilstein Archives with doi: 10.3762/bxiv.2020.43.v1 and is considered to be an early communication for feedback before peer review. Before citing this document, please check if a final, peer-reviewed version has been published in the Beilstein Journal of Organic Chemistry.

This document is not formatted, has not undergone copyediting or typesetting, and may contain errors, unsubstantiated scientific claims or preliminary data.

Preprint Title FERROCENYL-SUBSTITUTED TETRAHYDROTHIOPHENES VIA FORMAL (3+2)-CYCLOADDITION REACTIONS OF FERROCENYL THIOKETONES WITH DONOR-ACCEPTOR CYCLOPROPANES

Authors Grzegorz Mloston, Mateusz Kowalczyk, André U. Augustin, Peter G. Jones and Daniel B. Werz

Publication Date 06 Apr 2020

Article Type Full Research Paper

Supporting Information File 1 Mloston-M-398-SI-final for submission-04-04.docx; 2.0 MB

ORCID® iDs Grzegorz Mloston - <https://orcid.org/0000-0003-3631-3239>; Mateusz Kowalczyk - <https://orcid.org/0000-0002-8589-7704>

**FERROCENYL-SUBSTITUTED TETRAHYDROTHIOPHENES
VIA FORMAL (3+2)-CYCLOADDITION REACTIONS OF
FERROCENYL THIOKETONES WITH DONOR-ACCEPTOR
CYCLOPROPANES**

Grzegorz Mlostoń,^{a)}* Mateusz Kowalczyk,^{a,b)} André U. Augustin,^{c)} Peter G. Jones,^{d)}
and Daniel B. Werz^{c)}*

In Memoriam of Rolf Huisgen (München) 1920-2020

^{a)} University of Łódź, Department of Organic & Applied Chemistry, Tamka 12, 91-403
Łódź, Poland

^{b)} University of Łódź, PhD School of Exact Sciences

^{c)} Technische Universität Braunschweig, Institute of Organic Chemistry, Hagenring 30,
38106 Braunschweig, Germany

^{d)} Technische Universität Braunschweig, Institute of Inorganic and Analytical
Chemistry, Hagenring 30, 38106 Braunschweig, Germany

Email:


grzegorz.mloston@chemia.uni.lodz.pl; d.werz@tu-braunschweig.de

* Corresponding authors

Keywords:

ferrocenyl thioketones; donor-acceptor cyclopropanes; (3+2)-cycloaddition reactions;
sulfur heterocycles; tetrahydrothiophenes

Graphical abstract




Abstract

Ferrocenyl thioketones react with donor-acceptor cyclopropanes in dichloromethane at room temperature in the presence of catalytic amounts of $\text{Sc}(\text{OTf})_3$ yielding tetrahydrothiophene derivatives, products of formal (3+2)-cycloaddition reactions, in moderate to high yields. In all studied cases, dimethyl 2-arylcyclopropane dicarboxylates react with corresponding aryl ferrocenyl thioketones in a completely diastereoselective manner to form single products in which (C-2)-Ar and (C-5)-ferrocenyl groups are oriented in a *cis*-fashion. In contrast, the same cyclopropanes undergo reaction with alkyl ferrocenyl thioketones to form nearly equal amounts of both diastereoisomeric tetrahydrothiophenes. Low selectivity was also observed in the reaction of 2-phthalimide-derived cyclopropanes with ferrocenyl phenyl thioketone.

Introduction


Functionalized tetrahydrothiophenes constitute an important group of five-membered sulfur heterocycles; many of them, both chiral and achiral, with biotin as the best-known representative, form the key motif in numerous compounds of great practical importance [1,2]. The development of chemo- and diastereoselective syntheses for these compounds is thus a challenging problem. An elegant and highly efficient method for the construction of the tetrahydrothiophene ring is based on 1,3-dipolar cycloadditions of in-situ-generated thiocarbonyl *S*-methanides (thiocarbonyl ylides) with electron-deficient ethylenic dipolarophiles. This method was extensively

developed by Huisgen and his co-workers in the 1980s [3-5]. In the course of these studies, a non-orthodox stepwise mechanism of the 1,3-dipolar cycloaddition established by experiments performed with the sterically crowded thiocarbonyl *S*-methanide **1**, derived from 2,2,4,4-tetramethyl-3-thioxocyclobutanone and extremely electron-deficient ethylenes **2** such as (*E*)- and (*Z*)-dialkyl dicyanobutenoates (R = CO₂Me) [6], tetracyanoethylene (R = CN) [7] or (*E*)- and (*Z*)-1,2-bis(trifluoromethyl)ethylene-1,2-dicarbonitrile (R = CF₃) [8]. Both five-membered spiro-tetrahydrothiophenes **3** and seven-membered *S,N*-heterocycles (ketene imines) **4** were observed in the course of these reactions (Scheme 1). The latter products were trapped with suitable nucleophiles (R = CO₂Me) or even isolated and identified by means of spectroscopic methods (R = CF₃).



Scheme 1. Synthesis of spiro-tetrahydrothiophenes **3** via non-concerted (3+2)-cycloadditions of thiocarbonyl ylide **1** with electron-deficient ethylenes **2**; cyclic keteneimines **4** are also formed as products of formal (4+3)-cycloadditions.

In a recent work, an alternative, efficient and useful method for the synthesis of highly functionalized tetrahydrothiophenes of type **6** was reported [9] (Scheme 2). Under Lewis acid catalysis, formal (3+2)-cycloadditions of aromatic and cycloaliphatic thioketones (also thioesters) with donor-acceptor cyclopropanes **5** (D-A cyclopropanes) were realized.



Scheme 2. Formal (3+2)-cycloadditions of thioketones and (4+3)-cycloadditions of thiochalcones with donor-acceptor cyclopropanes **5** leading to tetrahydrothiophenes **6** and thiines **7**, respectively.


In contrast, thiochalcones (α,β -unsaturated aromatic thioketones) were shown to react under similar conditions with cyclopropanes **5** yielding exclusively seven-membered thiines **7** as products of the formal (4+3)-cycloaddition [10] (Scheme 2).

In a series of our recent publications, ferrocenyl/aryl and ferrocenyl/alkyl thioketones were demonstrated to be attractive substrates for the preparation of six- and five-membered sulfur heterocycles via (4+2)- and (3+2)-cycloadditions, respectively [11-14]. Notably, in contrast to aryl/alkyl thioketones (e.g. thioacetophenone), their ferrocenyl analogues of type **8** (e.g. ferrocenyl phenyl thioketone (**8a**), diferrocenyl thioketone (**8b**) and ferrocenyl methyl thioketone (**8c**)) are stable compounds at ambient conditions and can be used with no special precautions. In general, ferrocene has been considered as an 'exceptional compound' [15,16] and in our hands ferrocenyl-functionalized sulfur heterocycles, e.g. thiiranes and 1,3-dithiolanes, have found applications for the synthesis of compounds relevant for medicinal [17] and materials chemistry, and electrochemical studies [18].

In continuation of our studies on organic sulfur compounds and the mechanisms of their reactions, the main goal of the present work was the examination of the formal (3+2)-cycloaddition reactions of ferrocenyl-substituted thioketones **8** with D-A cyclopropanes **5**, aimed at the synthesis of hitherto unreported, ferrocenyl-substituted tetrahydrothiophene dicarboxylates (thiolanes) of type **9**.

Results and Discussion

In analogy to experiments described in our earlier publication [9], the test reaction was performed with dimethyl 2-phenylcyclopropane dicarboxylate (**5a**) and ferrocenyl phenyl thioketone (**8a**) in CH₂Cl₂ at room temperature using aluminium chloride (AlCl₃) as a catalyst. The reaction was monitored by TLC, and was shown to be complete after 1 h. The crude reaction mixture was examined by ¹H NMR, revealing the formation of a single product with characteristic signals of both CO₂Me groups located at 3.38 and 3.81 ppm. After chromatographic separation the expected tetrahydrothiophene **9a** was isolated in only 23% yield. As the next model substrate, the sterically crowded diferoceenyl thioketone (**8b**) was tested as a structural analogue of thiobenzophenone, which is widely applied in studies involving aromatic thioketones [3-5]. However, in contrast to **8a**, the reaction of **8b** with **5a** was unsuccessful; this observation prompted us to replace AlCl₃ by scandium triflate (Sc(OTf)₃), which is also known to be an efficient catalyst in various reactions of D-A cyclopropanes [9,10,19]. This time, the reaction was complete after 1 h and the expected 2,2-diferoceenyl-substituted tetrahydrothiophene **9b** was isolated chromatographically in about 28% yield (Scheme 3, Table 1). This experiment was successfully repeated, again using Sc(OTf)₃ instead of AlCl₃, in further experiments of ferrocenyl thioketones **8** with differently substituted cyclopropanes **5**. Again using Sc(OTf)₃, we repeated the experiment with **8a**, which this time led to the isolation of **9a** in an excellent yield of 98% (Table 1).



Ar = **a**: Ph; **b**: β-naphthyl;

c: 4-Me-C₆H₄;

d: 4-MeO-C₆H₄;

e: 4-Br-C₆H₄;

f: 4-CF₃-C₆H₄;

g: 1-phthalimid-1-yl

R = **a**: Ph; **b**: Fc;

c: β-naphthyl; **d**: Me;

e: *n*-Pr;

f: thien-2-yl;

g: fur-2-yl

Scheme 3. Formal (3+2)-cycloadditions of dimethyl 2-substituted cyclopropane-1,1-dicarboxylates **5a-g** with ferrocenyl thioketones **8a-g**, leading to dimethyl tetrahydrothiophene 3,3-dicarboxylates **9a-n** (Table 1).

Table 1.

Ferrocenyl-substituted tetrahydrothiophenes **9a-n** obtained in reactions of D-A cyclopropanes **5a-h** with ferrocenyl thioketones **8a-g** catalysed with Sc(OTf)₃.


Compound 9	Substituent Ar	Substituent R	Ratio of diastereoisomers	Yield of isolated products (%)
a	Ph	Ph	100 : 0	98
b	Ph	Fc	-	28
c	Ph	β-Naphthyl	100 : 0	65
d	Ph	Me	55 : 45	98
e	Ph	<i>n</i> -Pr	52 : 48	97
f	β-Naphthyl	Ph	100 : 0	30
g	β-Naphthyl	β-Naphthyl	100 : 0	31
h	4-Me-C ₆ H ₄	Ph	100 : 0	85
i	4-MeO-C ₆ H ₄	Ph	100 : 0	79
j	4-Br-C ₆ H ₄	Ph	100 : 0	93
k	4-CF ₃ -C ₆ H ₄	Ph	100 : 0	95
l	Ph	Thien-2-yl	100 : 0	58
m	Ph	Fur-2-yl	60 : 40	96
n	Phthalimid-1-yl	Ph	60 : 40	34

Fc = ferrocenyl

In analogy to **8a**, the similarly substituted ferrocenyl (β-naphthyl) thioketone (**8c**) reacted with **5a** in a diastereoselective manner yielding the expected **9c** in good yield (65%) as the sole isolated product. Notably, in all reactions performed with aryl-

substituted cyclopropanes **5a-f** and with thioketones **8a,c,f**, the desired tetrahydrothiophenes **9a,c,f-I** were formed with complete diastereoselectivity, leading to a single isomer. In order to establish the structure of the isomers, a single crystal obtained for compound **9c** was studied by X-ray diffraction analysis, which showed that the Ph(C-2) group and Fc(C-5) substituent are mutually *cis*-oriented (Figure 1). Tentatively, the same configuration was also attributed to all tetrahydrothiophenes **9a,f-I** that were formed as single isomers (Table 1).

However, the diastereoselectivity changed in reactions that were conducted with alkyl ferrocenyl thioketones **8d-e** with **5a**. Thus, the reaction with **8d** led to a 53:47 mixture of two isomeric products in nearly quantitative yield (98%). Subsequently, they were carefully separated by preparative thin layer chromatography (PTC) on silica using a mixture of petroleum ether and ethyl acetate as an eluent. The less polar fraction formed the major product and the slightly more polar one was isolated and identified as the minor isomer of **9d**. In the course of crystallization from hexane the less polar fraction gave single crystals suitable for the X-ray diffraction analysis which unambiguously confirmed that in this molecule the Ph(C-5) and Fc(C-2) groups are *trans*-oriented and for that reason, this isomer will be described as *trans*-**9d** (Figure 1).







Figure 1. Thermal ellipsoid plots of the molecular structures of *cis*-**9c** (above) and *trans*-**9d** (below) drawn using 50% probability displacement ellipsoids. *cis* and *trans* refer to the relative orientation of Ph and Fc groups).

Analogously, the reaction of ferrocenyl *n*-propyl thioketone (**8e**) with **5a** led to a 54:46 mixture of *trans*- and *cis*-isomers of **9e**, which were isolated in a total yield of 97% and identified without further separation. Moreover, a mixture of nearly equal amounts of isomeric *trans*-**9m** and *cis*-**9m** was also observed in the reaction of **5a** with ferrocenyl furan-2-yl thioketone (**8g**); the reaction of phthalimide-derived cyclopropane **5f** with thioketone **8a** led to a 4:1 mixture of both isomers *cis*- and *trans*-**9n**. Based on these observations it is difficult to explain the complete diastereoselectivity of tetrahydrothiophene formation observed in reactions of aryl ferrocenyl-substituted thioketones **8a,c,f** with cyclopropanes **5a-f** bearing aryl groups. Tentatively, a repulsive interaction of aryl groups rather than steric hindrance of the bulky ferrocenyl unit can be postulated. Remarkably, ferrocenyl fur-2-yl thioketone (**5g**) was an exception and delivered a 60:40 mixture of *trans*- and *cis*-**9m**.

The mechanistic interpretation of the efficient, formal (3+2)-cycloadditions of D-A cyclopropanes **5** with ferrocenyl thioketones **8** in the presence of a Lewis acid is based on the assumption that the coordination of the catalyst by two ester groups activates the cyclopropane ring and allows a nucleophilic attack of the C=S group on the benzylic position of the cyclopropane derivative (Scheme 4).



Scheme 4. Plausible mechanism for the formal (3+2)-cycloadditions of ferrocenyl thioketones **8** with D-A cyclopropanes **5**.

Subsequent ring-closure of the zwitterionic intermediate **10** leads to the formation of the tetrahydrothiophene derivative **9**. This process formally resembles the (3+2)-cycloadditions of thiocarbonyl *S*-methanides with an activated C=C double bond, which also lead to tetrahydrothiophenes [6-8]. Nevertheless, the key step involves formation of the reactive, zwitterionic intermediate **10**. It seems that repulsive interactions of aryl groups Ar (from cyclopropane **5**) and R (from thioketone **8**) control the diastereoselective ring-closure to the five-membered ring leading in these cases to the formation of *cis*-**9** (Ar to Fc) as a single isomer. A similar reaction pathway with a zwitterionic intermediate analogous to **10**, generated in the presence of a Lewis acid was proposed for the reaction of cycloaliphatic 3-thioxo-2,2,4,4-tetramethylcyclobutanone with D-A cyclopropanes [20].

Conclusions

The present study showed once more that ferrocenyl/aryl and ferrocenyl/alkyl thioketones **8** are versatile and useful building blocks for a simple and efficient

preparation of ferrocenyl-functionalized five-membered sulfur heterocycles. They were shown to react easily with donor-acceptor (D-A) cyclopropanes in the presence of scandium triflate $\text{Sc}(\text{OTf})_3$ as a catalyst, yielding highly functionalized tetrahydrothiophene derivatives of type **9**. These formal (3+2)-cycloaddition reactions occur via nucleophilic attack of the sulfur atom on the activated cyclopropane ring at the most reactive benzylic position. Formation of the five-membered ring occurs regioselectively and the expected tetrahydrothiophene-3,3-carboxylates are the products. The studied reactions display an interesting stereoselectivity and, in the case of 2,5-diaryl substituted products **9**, both aryl groups are exclusively located at the opposite sides of the ring plane. The described reactions supplement the recently reported synthetic applications of ferrocenyl thioketones [21], especially the well-known (3+2)- and (4+2)-cycloadditions leading to diverse sulfur heterocycles in which they played the role of reactive dipolarophiles (1,3-dipolar cycloaddition or the Huisgen reaction) [12-14,17] and dienophiles (*hetero*-Diels-Alder reaction) [11,18,22]. Because of the presence of the redox-active ferrocenyl unit, the tetrahydrothiophenes **9** described here could be of interest for further studies oriented towards new materials, biologically active compounds, or crystal engineering [15,16].

It should be also emphasized that the present study demonstrates the growing potential of donor-acceptor cyclopropanes [23-27] as unique building blocks for current organic synthesis and especially for efficient and highly stereoselective preparation of the relevant five-membered sulfur heterocycles derived from tetrahydrothiophene.

Experimental

General information: Solvents and chemicals were purchased and used as received without further purification. Products were purified by standard column chromatography on silica gel. Yields refer to analytically pure samples. NMR spectra were recorded with a Bruker Avance III 600 MHz instrument (^1H NMR: 600 MHz; ^{13}C NMR: 151 MHz). Chemical shifts are reported relative to solvent residual peaks (^1H NMR: $\delta = 7.26$ ppm [CHCl_3]; ^{13}C NMR: $\delta = 77.0$ ppm [CDCl_3]). IR spectra were recorded with a Cary 630 FTIR (*Agilent Technologies*) spectrometer (as film). Melting points were determined in capillaries with a Melt Temp II apparatus.

Starting materials: D-A cyclopropanes **5a-g** were obtained following the reported procedure [27]. Ferrocenyl thioketones **8a-g** were obtained by thionation of corresponding ferrocenyl ketones [28] by treatment with Lawesson's reagent [29]. Ferrocenyl β -naphthyl thioketone (**8b**) obtained from ferrocenyl(β -naphthyl) ketone [30] is reported for the first time (see SI).

General procedure: A solution of 0.3 mmol of the corresponding cyclopropane **5** in 5 mL of dichloromethane was stirred for 5 min. Then, 0.5 mmol of the corresponding ferrocenyl thioketone **8** and a catalytic amount (ca. 5 mg) of Sc(OTf)₃ was added to the stirred solution. The mixture was stirred at room temperature for 1 h. Progress of the reaction was monitored by TLC. The solvent was evaporated in vacuo and the crude mixtures were purified by flash chromatography using dichloromethane as the eluent. Analytically pure samples of tetrahydrothiophenes **9** were obtained by crystallization from petroleum ether or hexane with a small amount of dichloromethane.

Dimethyl 2-ferrocenyl-2,5-diphenyltetrahydrothiophene-3,3-dicarboxylate (cis-9a): Yield: 159 mg (98%). Red crystals; mp 192-193°C; ¹H NMR δ 2.65 (*dd*, $J_{H,H} = 13.9$ Hz, $J_{H,H} = 4.3$ Hz, 1H, HC(4)), 3.44 (*s*, 3H, OCH₃), 3.46 (*s*, 3H, OCH₃), 3.56 (*dd*, $J_{H,H} = 13.9$ Hz, $J_{H,H} = 12.7$ Hz, 1H, HC(4)), 3.51-3.60 (*m*, 1H, HC(Fc)), 4.00-4.02 (*m*, 1HC(Fc)), 4.07 (*s*, 5HC(Fc)), 4.27-4.29 (*m*, 1HC(Fc)), 4.68-4.70 (*m*, 1 HC(Fc)), 4.81 (*dd*, $J_{H,H} = 12.6$ Hz, $J_{H,H} = 4.3$ Hz, HC(5)), 7.31-7.39 (*m*, 2 arom. HC), 7.40-7.46 (*m*, 4 arom. HC), 7.62-7.65 (*m*, 2 arom. HC), 8.21-8.24 (*m*, 2 arom. HC); ¹³C NMR δ (C(4)-not found), 48.1, 48.4 (2OCH₃), 52.4 (C(5)), 67.8 (C(2)), 71.1 (C(3)), 68.7, 69.2, 69.9, 71.0, 73.7 (for 9 HC(Fc)), 97.0 (C(Fc)), 126.6, 127.1, 127.9, 128.0, 128.8, 129.0 (for 10 arom. HC), 138.9, 144.2 (2 arom. C), 169.0, 170.3 (2 C=O); IR ν 1737*brs* (2C=O), 1492*m*, 1444*m*, 1429*m*, 1258*m*, 1239*s*, 1073*m*, 814*m*, 760*m*, 697*vs*, 497*vs* cm⁻¹; Anal. calcd for C₃₀H₂₈FeO₄S (540.45): C 66.67, H 5.22, S 5.93; found: C 66.58, H 5.24, S 5.99.

Dimethyl 2,2-diferrocenyl-5-phenyl tetrahydrothiophene-3,3-dicarboxylate (9b): Yield: 54 mg (28%). Red crystals; mp 170°C (dec.); ¹H NMR δ 2.91 (*dd*, $J_{H,H} = 14.0$ Hz, $J_{H,H} = 6.2$ Hz, 1H, HC(4)), 3.36 (*s*, 3H, OCH₃), 3.41 (*dd*, $J_{H,H} = 15.8$ Hz, $J_{H,H} = 12.4$ Hz, 1H, HC(4)), 3.65 (*s*, 3H, OCH₃), 4.05-4.07 (*m*, 1H, HC(Fc)), 4.07-4.09 (*m*, 1H, HC(Fc)),

4.16-4.18 (*m*, 1H, HC(Fc)), 4.23-4.27 (*m*, 7H, 7HC(Fc)), 4.31 (*s*, 5H, HC(Fc)), 4.48-4.50 (*m*, 1H, HC(Fc)), 4.56-4.58 (*m*, 1H, HC(Fc)), 4.68-4.70 (*m*, 1H, HC(Fc)), 5.51 (*dd*, $J_{H,H} = 11.2$ Hz, $J_{H,H} = 6.3$ Hz, 1H, HC(5)), 7.34-7.37 (*m*, 1arom. HC), 7.44-7.48 (*m*, 2arom. HC), 7.78 (*m*, 2arom. HC); ^{13}C NMR δ 47.9 (C(4)), 49.5 (C(5)), 51.9, 52.6 (2OCH₃), 65.8, 66.2, 66.9, 67.6, 67.7, 69.6, 69.7, 70.4, 73.1 (for 18 HC(Fc)), 73.4, 94.3 (C(2) and C(3)), 100.0 (2C(Fc)), 127.4, 128.2, 128.6 (5 arom. HC), 141.2 (arom. C), 169.0, 169.1 (2 C=O); IR ν 1727brs (2C=O), 1431*m*, 1259*s*, 1164*s*, 1107*m*, 1000*m*, 818*s*, 760*m*, 696*s*, 479vs cm^{-1} ; Anal. calcd for C₃₄H₃₂Fe₂O₄S (648.37): C 62.98, H 4.97, S 4.94; found: C 62.68, H 4.93, S 4.88.

Dimethyl 2-ferrocenyl-5-phenyl-2-(naphth-2-yl)tetrahydrothiophene-3,3-dicarboxylate (*cis*-9c): Yield: 115 mg (65%). Yellow crystals; mp 210°-211°C; single crystals were obtained from hexane solution by slow evaporation at r.t. ^1H NMR δ 2.69 (*dd*, $J_{H,H} = 13.8$ Hz, $J_{H,H} = 4.3$ Hz, 1H, HC(4)), 3.41 (*s*, 3H, OCH₃), 3.47 (*s*, 3H, OCH₃), 3.57 (*s*, 1H, HC(Fc)), 3.62 (*t*, $J_{H,H} = 13.1$ Hz, 1H, CH), 4.01 (*s*, 1H, HC(Fc)), 4.09 (*s*, 5 HC(Fc)), 4.31 (*s*, 1H, HC(Fc)), 4.77 (*s*, 1H, HC(Fc)), 4.86 (*dd*, $J_{H,H} = 12.6$ Hz, $J_{H,H} = 4.3$ Hz, 1H, HC(5)), 7.36-7.40 (*m*, 1 arom., HC), 7.44-7.48 (*m*, 2 arom. HC), 7.51-7.54 (*m*, 2 arom. HC), 7.66 (*m*, 2 arom. HC), 7.87 (*d*, $J_{H,H} = 8.6$ Hz, 1 arom., HC), 7.88-7.92 (*m*, 1 arom., HC), 7.95-7.99 (*m*, 1 arom., HC), 8.36 (*d*, $J_{H,H} = 8.6$ Hz, 1arom., HC), 8.74 (*s*, 1 arom., HC); ^{13}C NMR δ 48.1 (C(5)), 48.5 (C(4)), 52.5, 52.6 (2OCH₃), 67.8, 68.7, 69.2, 69.9, 71.2 (for 9 HC(Fc)), 71.0, 73.5 (C(2) and C(3)) 97.4 (2 C(Fc)), 125.8, 125.9, 126.1, 127.2, 127.3, 127.9, 128.0, 128.1, 128.6, 128.8 (10 arom. HC), 132.1, 132.8, 138.8, 141.8 (4 arom. C), 168.9, 170.3 (2 C=O); IR ν 1738brs (2C=O), 1429*m*, 1239*s*, 1215*s*, 1170*m*, 1053*m*, 810*s*, 758*m*, 704*s*, 480vs cm^{-1} ; Anal. calcd for C₃₄H₃₀FeO₄S (590.51): C 69.15, H 5.12, S 5.43; found: C 67.16, H 5.01, S 5.47.

Dimethyl 2-ferrocenyl-2-phenyl-5-methyltetrahydrothiophene-3,3-dicarboxylate (9d). Obtained as a 55:45 mixture of isomers. *trans*- (major) and *cis*- (minor) isomers (Ph to Fc) were separated by PLC (silica, PE/ethyl acetate). Yields: *cis*-isomer – yellow crystals, 66 mg (more polar fraction, 45%); mp 148-150°C, *trans*-isomer – yellow crystals, 74 mg (less polar fraction, 55%); mp 126-128°C; single crystals of *trans*-9d were obtained from hexane/CH₂Cl₂ solution by slow evaporation at r.t. ^1H NMR (*cis*-9d) δ 2.31 (*s*, 3H, CH₃); 2.54 (*dd*, $J_{H,H} = 13.8$ Hz, $J_{H,H} = 5.3$ Hz, 1H, HC(4)); 3.22 (*dd*,

$J_{H,H} = 13.8$ Hz, $J_{H,H} = 12.3$ Hz, 1H, HC(4)); 3.46 (s, 3H, OCH₃); 3.80 (s, 3H, OCH₃); 4.09-4.10 (m, 1H, HC(Fc)); 4.12-4.13 (m, 1H, HC(Fc)); 4.22 (s, 5H, HCH(Fc)); 4.23-4.25 (m, 1H, HC(Fc)); 4.55-4.56 (m, 1H, HC(Fc)); 4.78 (dd, $J_{H,H} = 12.3$ Hz, $J_{H,H} = 5.3$ Hz, 1H, HC(5)); 7.30-7.34 (m, 1 arom. HC); 7.39-7.43 (m, 2 arom. HC); 7.54-7.58 (m, 2 arom. HC); ¹³C NMR (*cis-9d*) δ 25.9 (CH₃); 43.9 (C(4)); 47.9 (C(5)); 52.0, 52.6 (2OCH₃); 60.0 (C(2)); 68.1, 68.2, 68.8, 69.1, 70.7 (for 9 HC(Fc)); 70.6 (C(3)); 96.5 (C(Fc)); 127.5, 127.6, 128.7 (5 arom. HC); 140.0 (arom. C); 168.5, 169.7 (2C=O); IR ν 1731 *brvs* (2C=O); 1494*m*, 1453*m*, 1436*m*, 1248*vs*, 1207*m*, 1157*vs*, 1105*m*, 1038*s*, 829*m*, 766*s*, 702*vs* cm⁻¹; Anal. calcd for C₂₅H₂₆FeO₄S (478.38): C 62.77, H 5.48, S 6.70; found: C 62.69, H 5.52, S 6.63.

¹H NMR (*trans-9d*) δ 2.28 (s, 3H, CH₃); 2.79 (dd, $J_{H,H} = 14.1$ Hz, $J_{H,H} = 10.7$ Hz, 1H, HC(4)); 3.12 (dd, $J_{H,H} = 14.1$ Hz, $J_{H,H} = 7.1$ Hz, 1H, HC(4)); 3.47 (s, 3H, OCH₃); 3.66 (s, 3H, OCH₃); 4.19-4.21 (m, 2H, HC(Fc)); 4.23 (s, 5H, 5HC(Fc)); 4.40-4.42 (m, 1H, HC(Fc)); 4.56-4.57 (m, 1H, HC(Fc)); 5.28 (dd, $J_{H,H} = 10.7$ Hz, $J_{H,H} = 7.1$ Hz, 1H, HC(5)); 7.30-7.32 (m, 1 arom. HC); 7.39-7.42 (m, 2 arom. HC); 7.57-7.59 (m, 2 arom. HC); ¹³C NMR (*trans-9d*) δ 31.3 (CH₃); 47.1 (C(4)); 48.6 (C(5)); 52.2, 52.3 (2OCH₃); 60.1 (C(2)); 67.6 (C(3)); 68.5, 68.8, 69.1, 69.3, 71.1 (for 9HC(Fc)); 89.9 (C(Fc)); 127.3, 127.9, 128.6 (5 arom. HC); 142.6 (arom. C); 168.9, 169.5 (2C=O); IR ν 1737*vs*, 1720*vs* (2C=O); 1492*m*, 1453*m*, 1427*m*, 1258*vs*, 1220*s*, 1204*m*, 1106*m*, 1105*m*, 1023*m*, 993*m*, 829*m*, 766*s*, 703*vs* cm⁻¹; Anal. calcd for C₂₅H₂₆FeO₄S (478.38): C 62.77, H 5.48, S 6.70; found: C 62.70, H 5.46, S 6.59.

Dimethyl 2-ferrocenyl-5-phenyl-2-(thien-2-yl)tetrahydrothiophene-3,3-dicarboxylate (*trans-9l*): Yield: 95 mg (58%). Yellow crystals; mp 210°C (dec.); ¹H NMR δ 2.65 (dd, $J_{H,H} = 14.0$ Hz, $J_{H,H} = 4.5$ Hz, 1H, HC(4)), 3.43 (pseudo-*t*, $J_{H,H} = 13.9$ Hz, 1H, HC(4)), 3.48 (s, 3H, OCH₃), 3.52 (s, 3H, OCH₃), 4.10 (s, 5H, 5HC(Fc)), 4.14 (s, 2H, 2HC(Fc)), 4.30 (s, 1H, HC(Fc)), 4.68 (s, 1H, HC(Fc)), 5.02 (dd, $J_{H,H} = 13.6$ Hz, $J_{H,H} = 4.5$ Hz, 1H, HC(5)), 7.05-7.07 (m, 1 arom. HC), 7.21-7.23 (m, 1 arom. HC), 7.34-7.38 (m, 1 arom. HC), 7.42-7.45 (m, 2 arom., HC), 7.55-7.57 (m, 1 arom. HC), 7.60-7.63 (m, 2 arom. HC); ¹³C NMR δ 47.3 (C(4)), 49.0 (C(5)), 52.2, 52.5 (2OCH₃), 67.9, 69.0, 69.4, 70.4, 70.7 (for 9 HC(Fc)), 66.3, 74.2 (2 arom. C), 94.6 (1 C(Fc)), 123.4, 125.8, 126.8, 127.9, 128.0, 128.8 (for 8 arom. HC), 138.7, 150.3 (2 arom. C), 168.4, 169.5 (2C=O);

IR ν 1733brs (2C=O), 1427m, 1235s, 1146s, 1045m, 1032m, 818m, 766s, 691vs, 506m, 488s cm^{-1} ; Anal. calcd for $\text{C}_{28}\text{H}_{26}\text{FeO}_4\text{S}_2$ (546.48): HRMS (ESI⁺): calcd. for $[\text{C}_{28}\text{H}_{26}\text{FeO}_4\text{S}_2]^+$, 546.0621; found, 546.0629.

Supporting Information

CCDC-1992864 and CCDC-1992865 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk/structures>.

Supporting Information File 1

Experimental data for selected compounds **9**, details of the crystal structure determination, and the original ¹H and ¹³C NMR spectra for all products.

Acknowledgements

G. M. thanks Professor Wolfgang Weigand (Jena) (Alexander von Humboldt Foundation grant 'Institutspartnerschaft (Jena-Łódź 2018-2020), Professor Heinz Heimgartner (Zurich) and Dr. Katarzyna Urbaniak (Łódź) for stimulating discussions on the chemistry of ferrocenyl thioketones. D. B. W. thanks the European Union for an ERC Consolidator Grant (Project: GAINBYSTRAIN).

References

1. Benetti, S.; De Risi, C.; Gian P. Pollini, G. P.; Zanirato, V. *Chem. Rev.* **2012**, *112*, 2129–2163. doi: 10.1021/cr200298b
2. De Clercq, P. J. *Chem. Rev.* **1997**, *97*, 1755–1792. doi: org/10.1021/cr950073e
3. Huisgen, R.; Fulka, C.; Kalwinski, I.; Xingya, L.; Mloston, G.; Moran, J. R.; Pröbstl, A. *Bull. Soc. Chim. Belg.* **1984**, *93*, 511–532. doi: 10.1002/bscb.19840930701
4. Mlostoń, G.; Heimgartner, H. *Pol. J. Chem.* **2000**, *74*, 1503–1532. doi: not available.

5. Mlostoń, G.; Heimgartner, H. In: Padwa, A.; Pearson, W. H. (Editors) 1,3-Dipolar cycloaddition chemistry towards heterocycles and natural products. New York (NY): Wiley, 2002. pp. 315–360.
6. Huisgen, R.; Mlostoń, G.; Giera, H.; Langhals, E. *Tetrahedron* **2002**, *58*, 507–519. doi: 10.1016/S0040-4020(01)01147-4
7. Huisgen, R.; Mloston, G.; Langhals, E. *J. Org. Chem.* **1986**, *51*, 4085–4087. doi: 10.1021/jo00371a039
8. Huisgen, R.; Mlostoń, G.; Langhals, E.; Oshima, T. *Helv. Chim. Acta* **2002**, *85*, 2668–2685. doi: 10.1002/1522-2675(200209)85:9<2668::AID-HLCA2668>3.0.CO;2-3
9. Augustin, A. U.; Sensse, M.; Jones, P. G.; Werz, D. B. *Angew. Chem. Int. Ed.* **2017**, *56*, 14293–14296. doi: 10.1002/anie.201708346
10. Augustin, A. U.; Merz, J. L.; Jones, P. G.; Mlostoń, G.; Werz, D. B. *Org. Lett.* **2019**, *21*, 9405–9409. doi: 10.1021/acs.orglett.9b03623
11. Kowalski, K.; R. Karpowicz, R.; Mlostoń, G.; Miesel, D.; Hildebrandt, A.; Lang, H.; Czerwoniec, R.; Therrien, B. *Dalton Trans.*, **2015**, *44*, 6268–6276. doi: 10.1039/c5dt00246j
12. Hejmanowska, J.; Jasiński, M.; Mlostoń, G.; Albrecht Ł. *Eur. J. Org. Chem.*, **2017**, 950–954. doi: 10.1002/ejoc.201601307
13. Mlostoń, G.; Hamera, R.; Linden, A.; Heimgartner, H. *Beilstein J. Org. Chem.*, **2016**, *12*, 1421–1427. doi: 10.3762/bjoc.12.136
14. Mlostoń, G.; Hamera-Fałdyga, R.; Jeske, M.; Godziszewska, M.; Urbaniak, K.; Heimgartner, H. *J. Sulfur Chem.*, **2018**, *39*, 47–63. doi: 10.1080/17415993.2017.1363206
15. Astruc, D. *Eur. J. Inorg. Chem.* **2017**, 6–29. doi: org/10.1002/ejic.201600983
16. Heinze K., Lang H. (Guest eds), *Organometallics* (special issue), **2013B**, *32*, 5623–6146.
17. Mlostoń, G.; Hamera-Fałdyga, R.; Celeda, M.; Heimgartner, H. *Org. Biomol. Chem.* **2018**, *16*, 4350–4356. doi: 10.1039/c8ob01022f
18. Mloston, G.; Hamera-Fałdyga, R.; Heimgartner, H. *J. Sulfur Chem.*, **2018**, *38*, 267–278. doi: 10.1080/17415993.2017.1415339)
19. Petzold, M.; Jones, P. G.; Werz, D. B. *Angew. Chem. Int. Ed.* **2019**, *58*, 6225–6229. doi: 10.1002/anie.20181440

20. Augustin, A. U.; Busse, M.; Jones, P. G.; Werz, D. B. *Org. Lett.* **2018**, *20*, 820-823. doi: 10.1021/acs.orglett.7b03961
21. Cai, Z.; Liu, Ch.-X.; Wang, Q.; Gu, Q.; You, S.-L. *Nat. Commun.* **2019**, *10*, 4168. doi: org/10.1038/s41467-019-12181-x
22. Mlostoń, G.; Urbaniak, K.; Zimmer, R.; Reissig, H.-U.; Heimgartner, H. *ChemistrySelect*, **2018**, *3*, 11724–11728. doi: 10.1002/slct.201803
23. Reissig, H.-U.; Zimmer, R. *Chem. Rev.* **2003**, *103*, 1151–1196. doi: 10.1021/cr010016n
24. Schneider, T. F.; Kaschel, J.; Werz, D. B. *Angew. Chem. Int. Ed.* **2014**, *53*, 5504–5523. doi: 10.1002/anie.201309886
25. Cavitt, M. A.; Phun, L. H.; France, S. *Chem. Soc. Rev.* **2014**, *43*, 804–818. doi: org/10.1039/C3CS60238A
26. Werz, D. B.; Biju, A. T. *Angew. Chem. Int. Ed.* **2020**, *59*, 3385–3398. doi: 10.1002/anie.201909213
27. Kreft, A.; Lücht, A.; Grunenberg, J.; Jones, P. G.; Werz, D. B. *Angew. Chem. Int. Ed.* **2019**, *58*, 1955–1959. doi: 10.1002/anie.201812880
28. Mlostoń, G.; Hamera, R.; Heimgartner H., *Phosphorus Sulfur Silicon Rel. Elem.* **2015**, *190*, 2125–2133. doi: org/10.1080/10426507.2015.1071817
29. Mlostoń, G.; Pipiak, P.; Hamera-Fałdyga, R.; Heimgartner, H. *Beilstein J. Org. Chem.* **2016**, *12*, 716–724. doi: 10.3762/bjoc.12.71
30. C. D. Rodriguez-Cendejas, L. S. Liebeskind, E. Peña-Cabrera, *Archivoc* **2005**, (vi), 250–265. doi: 10.3998/ark.5550190.0006.621